

D3 10,000 + K~K2D3



Therapeutic Solutions

Clinical Applications

- Supports Bone Health by Promoting Carboxylation of Bone Proteins*
- Supports Cardiovascular Health by Affecting Arterial Calcium Deposits*
- Supports Healthy Blood Clotting*

*K2 Supreme, K2 Vitamin D3, and K2 Vitamin D3 5000 provide vitamin K2 as menaquinone-7 (MK-7), a highly bioavailable and bioactive form of K2. Historical use and numerous studies have demonstrated the efficacy of vitamin K supplements for bone and cardiovascular health.**

All Optimum Therapeutic Solutions Formulas Meet or Exceed cGMP Quality Standards

Discussion

Naturally occurring vitamin K is found as either K1 (phyloquinone), which is derived from food sources such as green leafy vegetables, or K2 (menaquinones). Menaquinones are designated as MK-n, where n denotes the length of the molecule's aliphatic side chain. Menaquinones are synthesized by bacteria and can be obtained from animal-based and fermented foods. Structural differences between K1 and K2 impact their bioavailability and bioactivity. Furthermore, among menaquinones, menaquinone-7 (MK-7), with its longer side chain, is very hydrophobic. Compared to K1, MK-7's physiochemical properties make it highly transportable by plasma lipoproteins, increase its extrahepatic (bones, arteries, etc.) availability, and produce its long half-life.^[1-3]

Absorption of K1 from food can be limited due to its membrane-bound nature and the individual consumer's digestive and absorptive variability. Moreover, adequate consumption of foods high in K2 can be challenging. Therefore, dietary supplementation is an important option. In addition, research suggests that higher levels of menaquinones are needed than were previously thought. Supplementary vitamin K can be found in three forms: synthetic K1; MK-4, which is structurally similar to K1; and natural, long-chain MK-7. Optimum Therapeutic Solutions provides MK-7 as Vitamk7™, a naturally derived and solvent-free vitamin K2 that has been obtained through a patent-granted biofermentation process of *Bacillus subtilis* natto cultures.*

MK-7 Bioavailability Increases Extrahepatic Tissue Utilization

Schurgers et al conducted human studies to compare the in vivo properties of orally administered K1 and MK-7. The results supported better bioavailability and utilization of MK-7. Expressed as AUC_{0-6h}, MK-7 demonstrated a six-fold better half-life, a seven- to eight-fold higher dose-response level, and a three times higher carboxylated to uncarboxylated osteocalcin ratio (cOC:ucOC[†]). Furthermore, on a molar basis, MK-7 is a three-to-four times more potent antidote for oral anticoagulation than is K1. Researchers note that, aside from sensitive individuals, "MK-7 supplements containing more than 50 mcg/d may interfere with oral anticoagulant treatment, whereas doses of at least 50 mcg are not likely to affect the INR value in a relevant way."^[2] Nonetheless, practitioners should closely monitor patients taking anticoagulants.*

While studies on the absorption and bioavailability of MK-4 at nutritional levels (i.e., doses of 500 mcg/d or lower) suggest less efficacy compared to longer-chain menaquinones at similar doses,^[4] this remains subject to debate. It is possible that rapid uptake of MK-4 could account for its observed lack of detection in serum after oral administration,^[5] but more studies are needed for clarification.*

Bone Benefits

Among the dietary factors critical to bone health, vitamin K has emerged as a key player. Vitamin K is believed to be necessary for bone mineralization. Through carboxylation, vitamin K activates osteocalcin, the protein needed to bind calcium to the mineral matrix in bone.^[6] Several studies have demonstrated the efficacy of MK-7 (e.g., doses of 45-90 mcg/d) to increase osteocalcin carboxylation and to increase the cOC:ucOC ratio. A high cOC:ucOC ratio is associated with bone health.^[1,2,4] A recent in vitro study also showed an osteogenic effect of MK-7 administration on human mesenchymal cell differentiation.^[6] In addition, the vitamin may protect bone integrity by reducing the synthesis of prostaglandin E2 or interleukin-6 by osteoclasts.^[7] Animal and human studies have demonstrated a significant beneficial effect of MK-7 supplementation on bone health.^[8-10] Vitamin K and vitamin D share some similar characteristics and are believed to act synergistically.^[11]

Cardiovascular and Other Health Benefits

Vitamin K benefits cardiovascular health by participating in the carboxylation of matrix GLA protein (MGP), a protein regarded to be the most potent inhibitor of arterial calcification. Researchers have demonstrated that supplementation with vitamin K reduces arterial calcium deposits^[1,3,12] and that long-term intake of long-chain menaquinones is inversely correlated with calcium accumulation in arteries.^[5]

Vitamin K has specific receptor binding sites that allow it to regulate gene activity.^[13] Besides its gene-mediating effects upon critical proteins, the vitamin can also bind with the steroid and xenobiotic receptors and influence their expression.^[14] In addition, vitamin K also demonstrates antioxidant activity^[15]; reduces levels of certain markers, such as acute phase reactants (e.g., C-reactive protein)^[16]; and participates in the induction of apoptosis.^[17]

†The cOC:ucOC ratio can be used as a determinant of vitamin K status.

***These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.**

Optimum Therapeutic Solutions
6420 N. MacArthur Blvd, Suite 100
Irving, TX 75039

D3 10,000 + K~K2D3

K2 Vitamin D3

Supplement Facts

Serving Size: 1 Capsule
Servings Per Container: 60

	Amount Per Serving	%Daily Value
Vitamin D3 (cholecalciferol)	10,000 IU	2500%
Vitamin K2 (as menaquinone-7)	45 mcg	56%

Other Ingredients: Microcrystalline cellulose, HPMC (capsule), vegetable stearic acid, vegetable magnesium stearate, and silica.

Directions

Swallow one capsule daily with water, preferably at mealtime, or as directed by your healthcare practitioner.

Consult your healthcare practitioner prior to use. Individuals taking other medication should discuss potential interactions with their healthcare practitioner. Consider total vitamin K intake (food and supplements) if you are taking blood-thinning medication. Present studies show that 45 mcg of MK-7 from Vitamk7™ daily is not likely to interfere with blood-thinning medicines. Do not use if tamper seal is damaged.

Does Not Contain

Wheat, gluten, yeast, soy protein, dairy products, fish, shellfish, peanuts, tree nuts, egg, ingredients derived from genetically modified organisms (GMOs), artificial colors, artificial sweeteners, or artificial preservatives.

References

1. Brugè F, Bacchetti T, Principi F, et al. Olive oil supplemented with menaquinone-7 significantly affects osteocalcin carboxylation. *Br J Nutr.* 2011 Oct;106(7):1058-62. [PMID: 21736837]
2. Schurgers LJ, Teunissen KJ, Hamulyák K, et al. Vitamin K-containing dietary supplements: comparison of synthetic vitamin K1 and natto-derived menaquinone-7. *Blood.* 2007 Apr 15;109(8):3279-83. [PMID: 17158229]
3. Beulens JW, Bots ML, Atsma F, et al. High dietary menaquinone intake is associated with reduced coronary calcification. *Atherosclerosis.* 2009 Apr;203(2):489-93. [PMID: 18722618]
4. Sato T, Schurgers LJ, Uenishi K. Comparison of menaquinone-4 and menaquinone-7 bioavailability in healthy women. *Nutr J.* 2012 Nov 12;11:93. [PMID: 23140417]
5. Schurgers LJ, Vermeer C. Differential lipoprotein transport pathways of K-vitamins in healthy subjects. *Biochim Biophys Acta.* 2002 Feb 15;1570(1):27-32. [PMID: 11960685]
6. Gigante A, Brugè F, Cecconi S, et al. Vitamin MK-7 enhances vitamin D3-induced osteogenesis in hMSCs: modulation of key effectors in mineralization and vascularization. *J Tissue Eng Regen Med.* 2012 Oct 29. [PMID: 23109511]
7. Weber P. Management of osteoporosis: is there a role for vitamin K? *Int J Vitam Nutr Res.* 1997;67(5):350-56. [PMID: 9350477]
8. Yamaguchi M, Taguchi H, Gao YH, et al. Effect of vitamin K2 (menaquinone-7) in fermented soybean (natto) on bone loss in ovariectomized rats. *J Bone Miner Metab.* 1999;17(1):23-29. [PMID: 10084398]
9. Knapen MH, Drummen NE, Smit E, et al. Three-year low-dose menaquinone-7 supplementation helps decrease bone loss in healthy postmenopausal women. *Osteoporos Int.* 2013 Sep;24(9):2499-507. [PMID: 23525894]
10. Kanellakis S, Moschonis G, Tenta R, et al. Changes in parameters of bone metabolism in postmenopausal women following a 12-month intervention period using dairy products enriched with calcium, vitamin D, and phyloquinone (vitamin K(1)) or menaquinone-7 (vitamin K (2)): the Postmenopausal Health Study II. *Calcif Tissue Int.* 2012 Apr;90(4):251-62. [PMID: 2239252]
11. Bolton-Smith C, McMurdo ME, Paterson CR, et al. Two-year randomized controlled trial of vitamin K1 (phyloquinone) and vitamin D3 plus calcium on the bone health of older women. *J Bone Miner Res.* 2007 Apr;22(4):509-19. [PMID: 17243866]
12. Geleijnse JM, Vermeer C, Grobbee DE, et al. Dietary intake of menaquinone is associated with a reduced risk of coronary heart disease: the Rotterdam Study. *J Nutr.* 2004 Nov;134(11):3100-05. [PMID: 15514282]
13. Igarashi M, Yogiashi Y, Mihara M, et al. Vitamin K induces osteoblast differentiation through pregnane X receptor-mediated transcriptional control of the Mx2 gene. *Mol Cell Biol.* 2007 Nov;27(22):7947-54. [PMID: 17875939]
14. Azuma K, Inoue S. Vitamin K function mediated by activation of steroid and xenobiotic receptor [in Japanese]. *Clin Calcium.* 2009 Dec;19(12):1770-8. [PMID: 19949268]
15. Vervoort LM, Ronden JE, Thijssen HH. The potent antioxidant activity of the vitamin K cycle in microsomal lipid peroxidation. *Biochem Pharmacol.* 1997 Oct 15;54(8):871-76. [PMID: 9354587]
16. Shea MK, Booth SL, Massaro JM, et al. Vitamin K and vitamin D status: associations with inflammatory markers in the Framingham Offspring Study. *Am J Epidemiol.* 2008 Feb 1;167(3):313-20. [PMID: 18006902]
17. Sada E, Abe Y, Ohba R, et al. Vitamin K2 modulates differentiation and apoptosis of both myeloid and erythroid lineages. *Eur J Haematol.* 2010 Dec;85(6):538-48. [PMID: 20887388]

***These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.**

Optimum Therapeutic Solutions
6420 N. MacArthur Blvd, Suite 100
Irving, TX 75039

DRS-233
REV. 08/02/16

Recommended Use: As a dietary supplement, take one softgel per day with a meal, or as directed by your health care practitioner.

Notice: Color, size or shape may appear different between lots.

✓ **MADE WITH NON-GMO INGREDIENTS**

✓ **DOES NOT CONTAIN GLUTEN**



Therapeutic Solutions

Co-Enzyme Q10

Optimum Absorption

Dietary Supplement

For Professional Use Only

60 SOFTGELS

Supplement Facts

Serving Size 1 softgel

Amount Per Serving	% Daily Value
Natural Coenzyme Q10 (Ubiquinone)	100 mg *

*Daily Value not established.

Other Ingredients: Medium chain triglycerides, sunflower lecithin, vitamin E, high gamma mixed tocopherols; gelatin, glycerine, purified water, annatto (color) (softgel ingredients).

Manufactured for: Optimum Therapeutic Solutions

6420 N. MacArthur Blvd., Suite 100, Irving, TX 75039

www.optimumhormonebalance.com

Report any adverse reactions to: 972-893-6068

STORE AT ROOM TEMPERATURE.

KEEP OUT OF REACH OF CHILDREN.

QEN100-PL-1

SUGGESTED USE: 1 soft gel capsule per day or as recommended by your health care professional. Formulated to be free of allergens derived from: Gluten, corn, yeast, artificial colors and flavors. If you are pregnant or nursing, consult your physician before taking this product. As with all dietary supplements, some individuals may not tolerate or may be allergic to the ingredients used. Please read the ingredient panel carefully prior to ingestion. Cease taking this product and consult your physician if you have negative reactions upon ingestion. KEEP CONTAINER TIGHTLY CLOSED. STORE AT ROOM TEMPERATURE. KEEP OUT OF REACH OF CHILDREN. This product was sealed for your protection. Do not use if outer looed neck seal or inner-seal is missing or damaged. 440.0040



Therapeutic Solutions

Opti-Omega 820

EPA/DHA (Molecularly Distilled)

Dietary Supplement
For Professional Use Only

120 SOFT GEL CAPSULES

Label ID L-HRN972-440120-C
Product #440120



Supplement Facts ^{v3}

Serving Size 1 Soft Gel Capsule
Servings Per Container 120

1 soft gel capsule contains	Amount Per Serving	% Daily Value
Calories	15	
Calories from fat	15	
Total fat	1.5 g	2%*
Cholesterol	<5 mg	<2%*
Total Omega-3s (as Triglycerides)	950 mg	**
EPA (Eicosapentaenoic Acid)	430 mg	**
DHA (Docosahexaenoic Acid)	390 mg	**
Omega-3s (additional)	130 mg	**

* Percent Daily Values are based on a 2,000 calorie diet.
** Daily Value not established

Ingredients: Highly Refined and Concentrated Omega-3 Fish Oil (anchovy), Capsule Shell (gelatin, glycerin, purified water), Natural Lemon/Lime Flavor, and a Proprietary Antioxidant Blend (consisting of natural tocopherols, rosemary extract, and ascorbyl palmitate).

Manufactured for: Optimum Therapeutic Solutions
6420 N. MacArthur Blvd, Suite 100, Irving, TX 75039
972-893-6068 • www.optimumhormonebalance.com

CUSTOMER HRN972

DATE 2-20-17

LABEL ID L-HRN972-440120-C

NOTICE This proof is a representation of your finished label. Artwork is subject to the output and settings of the monitors or printers used. Consequently colors on this proof should not be considered accurate. Your signature and return of this file ensures that you have given Ortho Molecular print approval after having proofread copy and examined the artwork presented on this proof. You are solely responsible for design; All elements of the text including: content, spelling, grammar and punctuation. Changes may delay your press date. Signed copy of this proof must be returned electronically or via fax before your order can be printed.

SIZE 8.815" x 3.313" BTL 400cc

COLORS PROCESS

FILENAME tL-HRN972-440120-C.ai

APPROVED CORRECTED PROOF REQUESTED



Client Signature _____ Date _____

Return Proofs to: Your Private Label Team
orthoprivatelabel@ompimail.com

1991 Duncan Place, Woodstock, IL 60098
ph: 815-206-6530 fax: 800-476-4664